arXiv:hep-ph/9409346v1 19 Sep 1994

Centre de Physique Théorique* - CNRS - Luminy, Case 907
F-13288 Marseille Cedex 9 - France

EXPANDING PROTONS SEEN BY HERA

Claude BOURRELY, Jacques SOFFER and Tai Tsun WU[1,2]

Abstract

We show that the rising total photoproduction cross section recently observed at HERA is entirely consistant with the impact-picture in the theory of expanding protons which was proposed more than twenty years ago. More accurate data, which is expected to be forthcoming soon, will give this prediction a stringent test.

Key-Words : Total photon cross section, impact-picture.

Number of figures : 2

August 1994
CPT-94/P.3060
CERN-TH/7387/94

anonymous ftp or gopher: cpt.univ-mrs.fr

---

* Unité Propre de Recherche 7061

[1] CERN - Geneva, Switzerland and Gordon McKay Laboratory, Harvard University, Cambridge, MA 02138, U.S.A.

[2] Work supported in part by the US Department of Energy under Grant N°. DE-FG02-84ER40158 and the Joint Service Electronics Program under Grant N00014-89-J-1023.

Recently at HERA, the $H1$ and ZEUS detectors have measured the photoproduction total cross section at an average γp center of mass energy of $180\ GeV$[1,2]. They confirm the previous and less accurate result of ZEUS[3] and indicate that this total cross section is rising. This rise has caused a lot of excitement and vastly different rates of increase have been proposed[4−9] which are based either on minijet contributions or on Regge - type analyses.

In this letter we will present a very simple approach to this increasing γp total cross section. Increases of hadron-hadron total cross sections have been theoreticaly predicted some years ago[10,11,12] and these predictions have been accurately verified by experiment[13]. The theoretical predictions were based on the multi-tower diagrams shown in fig.1a for pp and $\overline{p}p$ and fig.1b for $\pi^{\pm}p$ elastic scattering. The multi-tower diagrams for γp are shown in fig.1c. The similarity of the diagrams in fig.1b and fig.1c implies that the γp total cross section can be related approximately to those of $\pi^{\pm}p$. The major differences between these two figures are

i) The electromagnetic coupling e appears twice in fig.1c.

ii) The photon is its own antiparticle.

Because of those two differences it is natural to expect

$$\sigma_{\text{tot}}(\gamma p) = (\text{const.})\alpha \left(\sigma_{\text{tot}}(\pi^+ p) + \sigma_{\text{tot}}(\pi^- p)\right) \tag{1}$$

where α is the fine structure constant. This eq.(1) should hold for all center of mass energies above about $10 GeV$. The constant is to be determined from the low energy γp total cross section data[14]. Note that the contribution from ρ-exchange is cancelled in the sum, as it should be. For $\sigma_{\mathrm{tot}}(\pi^{\pm}p)$ we have used the early impact picture predictions[15] where a simple s dependence $s^{0.08}$ was first given and later extensively used by various authors. Using these results it is found that the constant in eq.(1) is 1/3 and thus the predicted γp total cross section is shown in fig.2. The HERA data is surely going to improve rapidly and we are eager to see a much more accurate determination of the γp cross section.

Thus we have encountered the expanding proton unambigously in two different experimental situations. In both cases, pp and γp, the next major increase in energy will come from LHC. In these cases the center of mass energy will reach 16 TeV for pp are more than 1 TeV for γp using electrons from LEP. Moreover using extracted π^+ and π^- beams from LHC, the πp cross sections could be measured up to about $\sqrt{s} = 120\ GeV$. This will provide a most stringent test of the accuracy of our relation (1).

Acknowledgments

Tai Tsun Wu wishes to thank Arthur Jordan for a helpful discussion and is grateful to John Ellis, Maurice Jacob, André Martin and many other members of the Theory Division for their hospitality at CERN.

References

[1] $H1$ Collaboration, T. Ahmed *et al.*, Phys. Lett. **B299** (1993) 374.

[2] ZEUS Collaboration, M. Derrick *et al.*, Preprint DESY 94-032 (march 1994).

[3] ZEUS Collaboration, M. Derrick *et al.*, Phys. Lett. **B293** (1992) 465.

[4] M. Drees and F. Halzen, Phys. Rev. Lett. **61** (1988) 275.

[5] R. Gandhi and I. Sarcevic, Phys. Rev. **D44** (1991) 10R.

[6] J.R. Forshaw and J.K. Storrow, Phys. Lett. **B268** (1991) 116.

[7] R.S. Fletcher, T.K. Gaisser and F. Halzen, Phys. Rev. **D45** (1992) 377.

[8] H. Abramowicz, E.M. Levin, A. Levy and U. Maor, Phys. Lett. **B269** (1991) 465.

[9] A. Donnachie and P.V. Landshoff, Phys. Lett. **B296** (1992) 227.

[10] H. Cheng and T.T. Wu, Phys. Rev. Lett. **24** (1970) 1456.

[11] C. Bourrely, J. Soffer and T.T. Wu, Phys. Rev. Lett. **54** (1985) 757.

[12] H. Cheng and T.T. Wu, Expanding protons : Scattering at high energies (MIT Press, Cambridge, MA, 1987).

[13] See for example N.A. Amos *et al.*, Phys. Lett. **B243** (1990) 158.

[14] D.O. Caldwell *et al.*, Phys. Rev. Lett. **40** (1978) 1222.

[15] H. Cheng, J.K. Walker and T.T. Wu, Phys. Lett. **B44** (1973) 97.

Figure Captions

Fig.1 Multi-tower diagrams for (a) pp and $\overline{p}p$, (b) $\pi^{\pm}p$ and (c) γp scattering.

Fig.2 γp total cross section as a function of the center of mass energy. The data are from refs. [1], [2] and [14] and the curve is from eq.(1).

This figure "fig1-1.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9409346v1

This figure "fig2-1.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9409346v1